

Rhythms of Recovery

Optimizing Sleep and Performance in Performing Artists

Joanna Fong-Isariyawongse, MD, FAAN, FAES

Associate Professor of Neurology, University of Pittsburgh

44th Annual PAMA International Symposium · From Practice to Performance: Occupational Health for Every Stage · July 2026

Sleep is a performance intervention, not an afterthought. It is behavioral, schedulable, dose-responsive, and it costs nothing. One behavior, one timing rule, one recovery window.

1. What each stage of sleep does for a performer

Stage	When it clusters	What it does	What the performer loses without it
N2	Across the night, roughly 50% of total sleep	Sleep spindles replay and consolidate motor sequences	Technique drilled yesterday does not stick. The passage you finally nailed is gone the next day.
N3 (slow wave)	First third of the night	Roughly 70% of daily growth hormone secretion, tissue repair, glycogen restoration, immune regulation, glymphatic clearance	Physical recovery and injury resilience. Cortisol stays elevated and the body never shifts into repair mode.
REM	Second half of the night	Memory consolidation and emotion regulation	Repertoire and choreography retention, plus composure under pressure. Cut sleep short and REM is lost first.

Practical consequence: cutting sleep at either end costs something different. Late bedtime costs slow wave sleep and physical recovery. Early alarm costs REM and emotional regulation on opening night.

2. Evidence at a glance

Sleep loss degrades performance

Outcome	Effect	Source
Skill accuracy and control	SMD -0.87	Kong et al., Front Physiol 2025
Explosive power	SMD -0.63	Kong et al., Front Physiol 2025
Aerobic endurance	SMD -0.59	Lopes et al., Sleep Med Rev 2023
Speed	SMD -0.52	Kong et al., Front Physiol 2025
Maximal force	SMD -0.35	Kong et al., Front Physiol 2025
Injury risk at under 8 hours per night	1.7x	Milewski et al., J Pediatr Orthop 2014; reviewed in Fox et al., Sports Med 2020
Long-term dysphonia at under 5 hours per night	1.6x	Cho et al., PLoS One 2017 (n=17,806)
Testosterone after 1 week at 5 hours	-10 to 15%	Leproult and Van Cauter, JAMA 2011
Insulin sensitivity	-10 to 30%	Endocrine literature, see references

Outcome	Effect	Source
Depression and anxiety risk, youth athletes	2 to 4x	Choi et al.; Orchard et al.

Skill accuracy and control carries the largest effect in this literature. For athletes it is one domain among five. For performing artists it is the entire performance surface: intonation, note accuracy, timing, pointe work, embouchure, breath control.

Sleep extension improves performance

Population	Intervention	Result
Collegiate basketball	+2 h/night, 5 to 7 weeks	Free throw accuracy +9%, three-point accuracy +9.2%, sprint time 4.3% faster (Mah et al., Sleep 2011)
Tennis	+2 h/night, 1 week	Serving accuracy +5%, daytime sleepiness improved
Ice hockey	10 h, single night	Cognitive performance +8%, fatigue reduced 23% (Alger et al., Sleep 2020)
Musicians (RCT, n=50)	+60 to 90 min/night, 7 nights	Musical memory +38%, rhythmic stability +4.5% (p<0.001), sightreading +7.1% (p<0.002), psychological flexibility increased (p<0.001) with controls unchanged (Pelletier, Music & Science 2025)
Napping (22 RCTs, n=291)	30 to 60 min, at least 1 h pre-performance	Physical performance SMD 0.99, cognitive SMD 0.69, fatigue SMD -0.76 (Mesas et al., BJSM 2023)

Note on the Pelletier memory figure: the 38% gain is a within-group improvement, and control participants also improved on the memorization task through practice effects. The sleep-attributable increment is smaller than the headline number. The rhythmic stability finding carries the strongest statistics.

3. Four behaviors to prescribe

- 1. Anchor wake time.** One wake time, held every day, including the morning after a show and including weekends. Bedtime will move. Wake time should not. This is the highest-yield single change because it stabilizes everything else.
- 2. Bank before demand.** Sleep debt cannot be repaid the night before, but it can be prepaid. Add 60 to 90 minutes per night in the week going into tech, a run, an audition, or a competition.
- 3. Use an off-ramp.** Performers come offstage aroused, not tired. Physiology needs 60 to 90 minutes to come down. Schedule that window rather than fighting it. No alcohol, no performance review.
- 4. Nap, then caffeinate.** 20 to 30 minutes, early afternoon, caffeine after and only if needed. Longer naps buy grogginess that is still being paid off at curtain.

4. Nap protocols

	Power nap: 20 to 30 min	Deep nap: 60 to 90 min
Use when	Well-rested, schedule is tight, close to curtain	Carrying sleep debt, touring, complex repertoire to consolidate
What it boosts	Alertness, mood, vigilance. Counteracts the post-lunch dip.	Sleep spindles and motor memory. Procedural consolidation of choreography and repertoire. Emotional reset.
Pre-curtain buffer	At least 30 minutes. Minimal sleep inertia.	At least 60 minutes, 2 hours ideal. Slow wave inertia impairs fine motor control.
Timing	13:00 to 15:00 post-lunch dip. Avoid after	Same window. Consider 100 to 200 mg

Power nap: 20 to 30 min	Deep nap: 60 to 90 min
16:00.	caffeine before the nap to blunt inertia.

Largest meta-analysis found the highest effect sizes in the 30 to under-60 minute range. When a full 90 minutes is not available, 30 to 45 minutes is the best practical compromise.

5. Production cycle strategy

The intervention changes depending on where the artist is in the cycle. Ask the question first: where are you in the production cycle?

Phase	Goal	Tactics
Rehearsal	Sleep banking	Extend sleep by 1 to 2 hours per night. Same wake time daily. Banking now creates the buffer later.
Tech week	Protect sleep	Guard the window. Draw on the bank. 20-minute afternoon nap.
The run	Optimize	60 to 90 minute wind-down after each show. 20-minute afternoon nap. No alcohol, no post-show review.
Dark days	Re-bank	Recovery sleep within 2 hours of the anchor wake time. Treat recovery as training.
Post-run	Full reset	Week 1 at 9 to 10 hours. Weeks 2 to 3 stabilize. Then bank for the next run.

Framework uses theater and dance. Musicians, singers, and touring artists can map the same principles onto season or tour structure.

6. Clinical workflow: screen, coach, escalate

Screen	Ask about frequency and impact. Three or more nights per week for three or more weeks means take action.
Coach	One specific behavior change: anchor wake time, morning light, or a wind-down ritual. Not all three.
Substances	Sedating is not restorative. Alcohol suppresses REM. Cannabis brings tolerance and withdrawal effects.
Guardrails	Caffeine under 400 mg per day, stopped 6 hours before bed. Naps 10 to 20 minutes or a full 90-minute cycle, ending 4 to 6 hours before bed.
Escalate	CBT-I is first-line for persistent insomnia. Digital CBT-I is a reasonable access route.

30-second script: *“Sleep is where today's rehearsal becomes tomorrow's performance. Let's pick the smallest schedule change that protects that this week.”*

Clinical aim: one behavior, one timing rule, one recovery window, all of it schedulable.

Selected references

Bagnall AD, Dorrian J, Fletcher A. Some vocal consequences of sleep deprivation and the possibility of fatigue proofing the voice with Voicecraft voice training. J Voice. 2011.

Cho JH, Guilleminault C, Joo YH, et al. A possible association between dysphonia and sleep duration. PLoS One. 2017.

Choi Y, Kim S, Park SH, et al. J Clin Med. 2025;14(2).

Craven J, McCartney D, Desbrow B, et al. Effects of acute sleep loss on physical performance: a systematic and meta-analytical review. Sports Med. 2022;52:2669-2690.

Facer-Childs E, Brandstaetter R. The impact of circadian phenotype and time since awakening on diurnal performance in athletes. Curr Biol. 2015.

Fox JL, Scanlan AT, Stanton R, Sargent C. Insufficient sleep in young athletes? Causes, consequences, and potential treatments. Sports Med. 2020;50(3):461-470.

Knaier R, et al. Diurnal variation in maximum endurance and strength performance. Med Sci Sports Exerc. 2022.

Kong Y, Yu B, Guan G, Wang Y, He H. Effects of sleep deprivation on sports performance and perceived exertion in athletes and non-athletes: a systematic review and meta-analysis. *Front Physiol.* 2025;16:1544286.

Leproult R, Van Cauter E. Effect of 1 week of sleep restriction on testosterone levels in young healthy men. *JAMA.* 2011;305(21):2173-2174.

Lopes TR, et al. Sleep restriction and aerobic endurance. *Sleep Med Rev.* 2023.

Mah CD, Mah KE, Kezirian EJ, Dement WC. The effects of sleep extension on the athletic performance of collegiate basketball players. *Sleep.* 2011;34(7):943-950.

Mesas AE, Nunez de Arenas-Arroyo S, Martinez-Vizcaino V, et al. Is daytime napping an effective strategy to improve sport-related performance and reduce fatigue? *Br J Sports Med.* 2023;57(7):417-426.

Milewski MD, Skaggs DL, Bishop GA, et al. Chronic lack of sleep is associated with increased sports injuries in adolescent athletes. *J Pediatr Orthop.* 2014;34(2):129-133.

Orchard F, Gregory AM, Gradisar M, Reynolds S. *J Child Psychol Psychiatry.* 2020;61(10):1126-1137.

Pelletier B. Effects of sleep extension on musical performance skills. *Music & Science.* 2025.

Questions and follow-up: fongjs@upmc.edu

Disclosure: speaker and consultant relationships with PepsiCo and Gatorade Sports Science Institute; speaker for Takeda Pharmaceuticals. These relationships do not influence the content or recommendations of this presentation.